

The MAGIC telescope

Ciro Bigongiari for the MAGIC collaboration* †

Padua University and INFN Padua, Italy

E-mail: ciro.bigongiari@pd.infn.it

MAGIC (Major Atmospheric Gamma Imaging Cherenkov telescope) is presently the largest ground-based gamma ray telescope. MAGIC has been taking data regularly since October 2004 at the Roque de los Muchachos Observatory on the island of La Palma. In this paper the MAGIC telescope status, its performances and some preliminary results on observed gamma ray sources are presented.

International Europhysics Conference on High Energy Physics

July 21st - 27th 2005

Lisboa, Portugal

*Speaker.

†Updated MAGIC members list at <http://wwwmagic.mppmu.mpg.de/collaboration/index.html>

1. Introduction

MAGIC (Major Atmospheric Gamma Imaging Cherenkov) telescope [1], is presently the largest imaging air Cherenkov telescope in operation. MAGIC has a 17 m diameter, $f/D = 1$, parabolic reflector covering a total surface of 234 m². The reflector dish is composed of 956 (0.495×0.495 m²) diamond milled aluminum mirrors [2]. The reflector shape is parabolic to minimize the time spread of Cherenkov light flashes on the focal plane. Aluminum mirrors were chosen instead of glass ones to reduce the weight of the reflecting surface and allow a fast slewing of the telescope. For the same reason the telescope frame is made of carbon fiber tubes. A very short slewing time is needed to catch the γ -ray prompt emission by GRBs. The reflected Cherenkov photons are recorded by a $3.5^\circ - 3.8^\circ$ FOV hexagonal camera in the telescope focal plane, composed by 397 0.1° FOV photomultiplier tubes, surrounded by 180 0.2° FOV PMTs. The PMTs have hemispherical windows and only 6 dynodes to minimize the time response width. The PMT photo conversion efficiency has been enhanced up to 30% and extended to the UV by coating the window with wavelength shifter [3]. The PMT signals are transferred via optical fibers [4] to the electronic room where they are split and sent to trigger and digitizing systems. The trigger decision is generated by a 2-level system using the signals of the 325 innermost PMTs. Only signals above an adjustable threshold are considered. The time coincidence within 6 nS of signals from 4 adjacent PMTs is required [5]. The analog signals are continuously digitized by 8 bit 300 MHz Flash ADCs. If the trigger condition is fulfilled the signals stored in FADC ring buffers are written to a FIFO buffer and saved by the DAQ.

2. MAGIC performances

The MAGIC construction was completed in Fall 2003 at the Roque de Los Muchachos observatory on the Canary island of La Palma, (28.75° N, 17.90° W, 2200 m above sea level). The commissioning phase took one year and in Fall 2004 MAGIC started to take data continuously on γ -ray sources. The first regular observation cycle started in April 2005. All the technical innovations implemented in the MAGIC construction [6] are now working without major problems and most of the telescope parameters are well within the design specifications [7]. According to detailed simulation of atmospheric showers and detector response the trigger threshold is around 60 GeV [8]. Presently the analysis threshold is about $E_{Th} = 100$ GeV as the γ /hadron discrimination becomes more and more difficult at lower energies. There is still room for lowering both trigger and analysis thresholds with present hardware fully exploiting second level trigger capabilities and improving the analysis technique. MAGIC integral flux sensitivity has been calculated from MC simulation and results in about 5% of Φ_{Crab} at $E > 100$ GeV and 2% of Φ_{Crab} at $E > 1$ TeV [8]. The angular resolution has been estimated applying the DISP method to Crab data and results in about 0.1° for γ -ray events with $E > 200$ GeV [9]. The accuracy in point-source position determination improves as the square root of the number of collected events and is ultimately limited by tracking accuracy ($\simeq 0.01^\circ$) [10]. The energy resolution has been estimated from MC data and results in $\Delta E/E \simeq 30\%$ at $E = 100$ GeV and $\Delta E/E \simeq 20\%$ for $E > 1$ TeV [11].

3. Gamma-ray source observations

Until Nov 2005 MAGIC has observed nearly 50 γ -ray source candidates. Emission of γ -rays has been detected by 8 of them, 4 galactic and 4 extragalactic.

The first and most observed source is the Crab Nebula. The Crab spectrum between some tens of GeV and some hundreds of GeV is astrophysically very interesting because the inverse Compton peak is expected to be close to 100 GeV and the cut-off of the pulsed γ -emission between 10 and 100 GeV. MAGIC has measured the Crab spectrum down to 100 GeV [11] for the first time, definitely heading for the inverse Compton peak. MAGIC detected γ -ray emission from the




Figure 1: Observed differential spectra of HESS J1813-178 and HESS J1834-087.




Figure 2: Observed differential spectrum of the source at galactic center.

SNRs, HESS J1813-178 and HESS J1834-087 [14], recently discovered by HESS [12]. MAGIC can observe these sources only at zenith angle around $40^\circ - 50^\circ$. This implies a higher energy threshold $E_{Th} = 300$ GeV but also a larger effective area. The spectra measured by MAGIC are in good agreement with HESS ones (Figure 1). The emission of VHE γ -rays from a source at galactic center discovered by Cangaroo and recently confirmed by HESS has been observed by MAGIC too [15]. Within errors the flux measured by MAGIC is compatible with the hard spectrum measured by HESS (Figure 2). The source position measured by MAGIC is compatible with both Sgr A* and Sgr A East.

MAGIC detected γ -rays from three known extragalactic sources, Mrk 421, Mrk 501 and 1ES 1959+650 and discovered γ -ray emission from another AGN, 1ES 1218+304, which is currently the second most distant gamma-ray source. Mrk 421 spectrum has been measured down to 100 GeV [13] for the first time. Flux variations of a factor 2 on an hourly time scale were observed with no evidence of spectral index variations. MAGIC detected an intense flare ($\Phi \simeq 4\dot{\Phi}_{Crab}$) from Mrk 501 on June 30th, 2005, while monitoring this source and alerted the astronomical community (IAU circular 8562). Very large flux variations were observed on short time scale. There are hints of spectral hardening when the source is in high flux state. The AGN 1ES 1959+650 spectrum has been measured by MAGIC down to 200 GeV when it was in low flux state [16]. This observation is characterized by lack of strong time variability and the presence of a significant VHE γ -ray emission in the absence of strong X-ray or optical emission. MAGIC discovered a new γ -ray source, the AGN 1ES 1218+304, and measured its spectrum between 100 GeV and 1 TeV. The measurement of VHE γ -ray spectra of distant AGNs, like 1ES 1218+304 can be used to evaluate the Metagalactic Radiation Field.

4. Conclusions and outlook

MAGIC has been taking data regularly since Fall 2004. About 50 possible γ -ray sources have been observed so far and γ -ray emission has been detected by 8 of them. MAGIC confirmed γ -ray emission from two sources recently discovered by HESS and detected γ -rays from 1ES 1218+304 at $z = 0.182$ for the first time. MAGIC is able to measure the emission spectrum of sources which culminate close to zenith down to 100 GeV. This limit will be lowered further by improving the data analysis technique, adopting new 2.5 GHz FADCs and building a second telescope. The construction of the second telescope, already started, will be completed by the end of 2007.

References

- [1] J. A. Barrio et al., *The MAGIC Telescope Design Report, MPI Report MPI-PhE/98-5*
- [2] C. Bigongiari et al., *The MAGIC Telescope Reflecting Surface NIM A 518 (2004) 193*
- [3] D. Paneque et al., *A method to enhance the sensitivity of photomultipliers for Air Cherenkov Telescopes by applying a lacquer that scatters light NIM A 518 (2004) 619*
- [4] D. Paneque et al., *Analogue signal transmission by an optical fiber system for the camera of the MAGIC telescope in proceedings of 28th International Cosmic Ray Conference (2003) 2927*
- [5] M. Meucci et al., *The Trigger System of the MAGIC Telescope: On-Line Selection Strategies for Cherenkov Telescopes, NIM A 518 (2004) 554*
- [6] R. Mirzoyan et al., *Technical Innovations for the MAGIC Project, in proceedings of 28th International Cosmic Ray Conference (2003) 2963*
- [7] J. Cortina et al., *Technical Performance of the MAGIC Telescope, in proceedings of 29th International Cosmic Ray Conference (2005) [astro-ph/0508274]*
- [8] P. Majumdar et al., *Monte Carlo simulation for the MAGIC telescope, in proceedings of 29th International Cosmic Ray Conference (2005) [astro-ph/0508244]*
- [9] E. Domingo Santamaria et al., *DISP analysis method for point-like off-axis and/or extended sources in proceedings of 29th International Cosmic Ray Conference (2005) [astro-ph/0508244]*
- [10] B. Riegel et al., *A tracking monitor for the MAGIC telescope in proceedings of 29th International Cosmic Ray Conference (2005) [astro-ph/0508274]*
- [11] R. Wagner et al., *Observations of the Crab nebula with the MAGIC telescope, in proceedings of 29th International Cosmic Ray Conference (2005) [astro-ph/0508244]*
- [12] F. Aharonian et al., *A new population of very high energy gamma-ray sources in the Milky Way Science 307 (2005) 1938 [astro-ph/0504380]*
- [13] D. Mazin et al., *Observations of Mrk 421 with the MAGIC Telescope in proceedings of 29th International Cosmic Ray Conference (2005) [astro-ph/0508244]*
- [14] H. Bartko et al., *Observations of VHE gamma radiation from HESS J1813-178 with the MAGIC telescope in proceedings of 29th International Cosmic Ray Conference (2005) [astro-ph/0508244]*
- [15] H. Bartko et al., *Search for Gamma Rays from the Galactic Center with the MAGIC telescope in proceedings of 29th International Cosmic Ray Conference (2005) [astro-ph/0508244]*
- [16] N. Tonello et al., *Observation of the AGN 1ES1959+650 with the MAGIC telescope in proceedings of 9th International Conference on Topics in AstroParticle and Underground Physics (2005)*